

# Medications Containing Aspirin

If you take any of the following aspirin-containing medications, stop taking the medicine 7 days prior to your procedure.

4-Way Cold Tablets	Disalcid	Omega-3 fatty acids, Fish Oil
Acuprin	Doan's Pills	Oxaprozin (Daypro)
Advil	Dolobid ( Diflunisal)	Pennsaid
Aggrastat (Tirofiban)	Dristan	Percodan
Aggrenox	Easprin	Persantine
Aleve	Edoxaban (Savayasa)	Piroxicam
Alka Seltzer (all varieties)	Ecotrin (all varieties)	Plavix (Clopidogrel)
Amytal with ASA	Effient (Prasugrel)	Pletal (Cilostazol)
Anacin (all varieties)	Eliquis (Apixaban)	Pradaxa (Dabigatran)
Ansaid (Flurbiprofen)	Empirin	Prexige (Lumiracoxib)
Anturane	Enoxaparin	Relafen (Nabumetone)
Arthritis Pain Formula (Whitehall)	Eptifibatide	ReoPro (Abciximab)
Arthropan	Equagesic	Rhumalgan
ASA (all varieties)	Etodolac (Ultradol)	Robaxisal
Ascaphen	Exedrin	Saleto
Ascodeen	Feldene	Salocol
Ascriptin	Fiorinal (all varieties)	Saw Palmetto
Aspergum	Garlic	Soma Compound
Aspirin	Ginko Bilboba	Stanback Powder
Axotal	Ginseng	Sulfinpyrazone
BA-C tablets	Goody's Powder (all varieties)	Sulindac (Clinoril)
Back-Quell	Halfprin 81 EC	Supac
Ban-O-Pain	Ibuprofen	Surgam
Bayer (all varieties)	Indomethacin (Indocid)	Synalgos DC
BC Powder (all varieties)	Integrelin (Eptifibatide)	Talwin
BiAct Cold	Jantoven	Tiaprofenic Acid (Surgam)
Brilinta (Ticagrelor)	Ketoprofen (Orudis)	Ticagrelor
Bufferin (all varieties)	Licofelone	Tenol Plus
Cama Arthritis Pain Reliever	Lortab ASA	Ticlid (Ticlopidine)
Cardiopirin	Lumiracoxib	Tolectin (Tolmetin)
Cataflam	Meloxicam (Mobic)	Trigesic
Celebrex (Celecoxib)	Methyl Salicylate (oil of wintergreen)	Tri-Pain Caplets
Cephalgesic	Midol (all varieties)	Ursinus Inlay Tabs
Cilostazol	Mobic (Meloxicam)	Vioxx (Rofecoxib)
Coricidin	Motrin	Vitamin C
Coumadin (Warfarin)	Multivitamin	Vitamin E
Dabigatran	Nabumetone	Voltaren, Voltarol
Damason-P	Naprosyn, Naproxyn Sodium, (Anaprox)	Warfarin
Diclofenac	Nimesulide (Eskafam, Mesulid)	Wesprin Buffered
Dicloflex, Difen, Difene	Norgesic Forte	Xarelto (Rivaroxaban)
Dipyramidole		Zorprin

Unless you have been told otherwise, do not eat or drink anything (not even gum, mints, or water) after midnight the night before surgery except medications for blood pressure, heart conditions, Parkinson's and seizures. Take these medications with only a tiny sip of water – enough to swallow the pill.