Prostate Biopsy Post-Procedure Instructions



Following the instructions below will contribute to a successful recovery.

Diet

You may return to your normal diet immediately.

Activities

Avoid strenuous activity for the first 24 hours. You may return to your normal activity the day after the biopsy.

Medication

- It is important to take the antibiotic (Cipro) that you were given for the biopsy the evening of your biopsy.
- You may take Tylenol (acetaminophen) if needed for pain or discomfort
- Do NOT take aspirin or any blood thinners like Coumandin, Plavix, Eliquis, Brilinta, or Pradaxa for 48 hours after your biopsy.
- Other than blood thinners, you should resume your regular medications.

Call your urologist if you experience any of these problems:

- Fevers over 101.5° F
- Heavy bleeding or clots
- Inability to urinate

Results

It will take approximately one week to get the results from your biopsy. If you have not heard from your doctor within two weeks after the biopsy, please call.

Clinic Locations

Medical Center

Pasteur Plaza 7909 Fredericksburg, Ste. 120-210 San Antonio, TX 78229 (210) 614-4544

Mission Trail

Mission Trail Medical Plaza 3327 Research Plaza, Ste. 403 San Antonio, TX 78235 (210) 337-6228 Metropolitan Prof. Bldg. 1303 McCullough, Ste. 270 San Antonio, TX 78212 (210) 474-7020

Downtown

Boerne

Methodist Med. Center 134 Menger Springs Rd., Ste. 1210 Boerne, TX 78006 (210) 614-4544

Northeast

Northeast Methodist Plaza 12709 Toepperwein, Ste. 206 San Antonio, TX 78233 (210) 564-8000

Westover Hills

Christus Medical Plaza 1 11212 Hwy. 151, Ste. 180 San Antonio, TX 78251 (210) 521-7333

Stone Oak

18915 Meisner Drive San Antonio, TX 78258 (210) 499-5158

Reyna Cancer Center

7718 Louis Pasteur Ct. San Antonio, TX 78229 (210) 477-9060

REV. 10/20/2017